

In Vitro

JMI Laboratories, North Liberty, Iowa, USA; Pfizer Inc., New York, NY, USA

Indicazioni

- Mafegia i a e e a if ga, age age ig he f ga, G 1 e . The d g f a age i (f a age i) i e e ig Pha e 3 c, i ca, de e, a f a a a f ca dide a/ i a i e ca didia i (NCT05421858).
- Mafegia ha i o, de n ed e in vitro acii agai . *Candida* ., e ce *C. krusei* (e a ed *Pichia kudriavzevii*), *Aspergillus* ., a d d i a e .
- Thi d de e n ed he in vitro acii f a age i a d ca a a agai . a age ca, ecio a fi f e , e ca e ad ea a d d i a e f n he SENTRY S ei, a ce P a g a n

Re

- Mafegia de n ed e in vitro acii agai . i f e e , e ca e ad ea . e hibi i g e e a ed MIC a, e . e h e d g c, a . e :
 - Candida fermentati* (MIC_{50/90}, 0.008/0.06 g/L),
 - C. guilliermondii* (MIC_{50/90}, 0.008/0.016 g/L),
 - C. quercitrusa* (MIC, 0.016 g/L), *Saprochaete clavate* (f a a Geotrichum clavatum; MIC_{50/90}, 0.03/0.06 g/L),